arXiv:1403.7029v1 [math.DG] 27 Mar 2014

On regular algebraic surfaces of $\mathbb{R}^3$ with constant mean curvature

J. Lucas M. Barbosa and Manfredo P. do Carmo *

Abstract

We consider regular surfaces M that are given as the zeros of a polynomial function $p : \mathbb{R}^3 \to \mathbb{R}$, where the gradient of p vanishes nowhere. We assume that M has non-zero mean curvature and prove that there exist only two examples of such surfaces, namely the sphere and the circular cylinder.

1 Introduction

An algebraic set in $\mathbb{R}^3$ will be the set

$$M = \{(x, y, z) \in \mathbb{R}^3;\ p(x, y, z) = 0\}$$

of zeros of a polynomial function $p : \mathbb{R}^3 \to R$. An algebraic set is regular if the gradient vector $\nabla p = (p_x, p_y, p_z)$ vanishes nowhere in M; here p_x, p_y and p_z denotes the derivative of p with respect to x, y or z respectively.

The condiction of regularity is essential in our case. It allows to parametrize the set M locally by differentiable functions $x(u, v)$, $y(u, v)$, $z(u, v)$ (not necessarily polynomials), so that M becomes a regular surface in the sense of differential geometry (see [2] chapter 2 section 2.2, in particular Proposition 2); here (u, v) are coordinates in an open set of $\mathbb{R}^2$.

Since M is a closed set in $\mathbb{R}^3$, it is a complete surface. In addition, being a regular surface, it is properly embedded, i. e., the limit set of M (if any)

*Both authors are partially supported by CNPq, Brazil

does not belong to M (cf. [15], chapter IV, A.1 p. 113). In particular, regular algebraic surfaces are locally graphs over their tangent planes,

From now on, M will denote a regular algebraic surface in $\mathbb{R}^3$. Due to the regularity condition, one can define on M the basic objects of Differential Geometry of surfaces and pose some differential-algebraic questions within this algebraic category.

For instance, in the last 60 years (namely after the seminal work [4] of Heinz Hopf in 1951), many questions have been worked out on differentiable. surfaces of non-zero constant mean curvature H. See also [5].

In our case, we have two examples of algebraic regular surfaces that have non-zero constant mean curvature, namely,

(1) Spheres, $(x-x_0)^2+(y-y_0)^2+(z-z_0)^2 = r^2$ with center $(x_0, y_0, z_0) \in \mathbb{R}^3$ and radius $r = 1/H$.

(2) Circular right cylinders, $(x - x_0)^2 + (y - y_0)^2 = r^2$, whose basis is a circle in the plane xy with center (x_0, y_0) and whose axis is a straight line passing through the center and parallel to the z axis.

A first natural question is: Are there further examples?

The first time we heard about this question was in a preprint of Oscar Perdomo (recently published in [13]) where he proves that for polynomials of degree three there are no such surfaces.

In this note, we prove the following general result:

Theorem: *Let M be a regular algebraic surface in $\mathbb{R}^3$. Assume that it has constant mean curvature $H \neq 0$. Then M is a sphere or a right circular cylinder.*

Acknowledgements. We want to thank Fernando Codá Marques for a crucial observation, Oscar Perdomo for having written [13] and Karl Otto Stöhr for his help in our first attempts to solve the problem.

2 Preliminaries

We first observe that, in the compact case, this theorem follows immediately from Alexandrov's well-known result: *An embedded compact surface in $\mathbb{R}^3$ with constant mean curvature is isometric to a sphere.*

The second observation is that the total curvature of an algebraic surface is finite. This was first proved by Osserman [11] in the case that the surface is an immersion parametrized by polynomials in two variables.

Here we give a proof of the finiteness of the total curvature for a (implicitly defined) regular algebraic surface M.

Proposition 1**:** *Let M be a regular algebraic surface in $\mathbb{R}^3$. Assume that it is complete non-compact. Then its total curvature is finite. More precisely,*

$$\int_M |K|\, dM \leq 4\pi C(d),$$

where K is the Gaussian curvature of M and $C(d)$ is a constant that depends only on the degree d of the polynomial defining M.

Proof. Let $g : M \to S^2(1)$ be the Gauss map of M. It is well known the Gauss curvature $K = \det(dg)$, where dg is the differential of the map g. Of course the result is true if M is either a plane or a circular cylinder. So, we ruled out these two cases from our proof. Let M^* be the set of points in M where $K \neq 0$. Then, restricted to M^*, g is a local diffeomorphism; that is, given $q \in g(M^*)$ and $m_\alpha \in \{g^{-1}(q)\}$, α belonging to a set of indices A, there exist neighborhoods $\mathcal{U}$ of q and $\mathcal{V}_\alpha$ of m_α such that, for each α, g maps V_α diffeomorphically onto $\mathcal{U}$. In fact, g restrict to M^* is a covering of $N(M^*)$ without ramification points. Since

$$\int_M |K| dM = \int_{M^*} |K| dM$$

the theorem is proved if we show that the mentioned covering has only a finite number of leaves.

Now, fix a plane P passing through the origin of $\mathbb{R}^3$ and in P fix an orthonormal basis $\{e_1, e_2\}$. Then, there exists a point $m \in M$ such that, up to translations, $P = T_m(M)$ if and only if

$$\frac{\nabla p}{|\nabla p|}(m) \perp P$$

what is equivalent to the system of equations

$$\begin{cases} \langle \frac{\nabla p}{|\nabla p|}(m), e_1 \rangle = 0 \\ \langle \frac{\nabla p}{|\nabla p|}(m), e_2 \rangle = 0 \end{cases} \tag{1}$$

Set $e_1 = \sum a_i U_i$ and $e_2 = \sum b_i U_i$ where $U_1 = (1,0,0)$, $U_2 = (0,1,0)$, $U_3 = (0,0,1)$ is the canonical basis of $\mathbb{R}^3$. Then, since $|\nabla p| \neq 0$, then (1) takes the form

$$\begin{cases} p_x a_1 + p_y a_2 + p_z a_3 = 0 \\ p_x b_1 + p_y b_2 + p_z b_3 = 0 \end{cases} \tag{2}$$

where the a_i, b_i, $i = 1,2,3$ are real numbers.

The equations in (2) describe algebraic surfaces, Σ_1 and Σ_2, which correspond to the coefficients a_i and b_i respectively, and whose degrees are $\leq (d-1)$, where d is the degree of p.

The surfaces Σ_1 and Σ_2, together with the original surface M determines points $m \in M$ as follows:

Σ_j interset M, $j = 1,2$, in a curve C_j. If the intersection $C_1 \cap C_2$ contains a point $m \in M^*$, since $K(m) \neq 0$, such intersection is unique in a neighborhood of m. By Bezout's theorem the total number of intersections is bounded above by $(d-1)^2$, as we wished. This proves the proposition.

Since $\int_M |K|\, dM$ is finite, it follows from a theorem of Huber [6] that the surface M of Propositon 1 is finitely connected, i.e., it is a compact surface with a finite number of ends.

The proof of our Theorem uses in a crucial way the structure theory for embedded, complete finitely connected surfaces with non-zero constant mean curvature developed by Korevaar, Kusner and Solomon in [7] after some preliminary work by Meeks [9]. The statement that we need from these papers is as follows:

Theorem A: ([9] and [7].) *Let M be a complete, non-compact, properly ebedded surface in $\mathbb{R}^3$ with non-zero constante mean curvature. Assume that M is finitely connected. Then, the ends of M are cylindrically bounded. Furthermore, for each end E of M, there exists a Delaunay surface $\Sigma \subset \mathbb{R}^3$ such that E and Σ can be expressed as cylindrical graphs ρ_E and ρ_Σ so that, near infinity, $|\rho_E - \rho_\Sigma| < Ce^{-\lambda x}$ where $C \geq 0$ and $\lambda > 0$ are constants.*

Remark: The first assertion in Theorem A comes from [9]. The final assertion is from [7], theorem 5.18.

3 Proof of the Theorem

We can assume that M is complete and non-compact; otherwise it is a sphere. Thus, by Proposition 1, M has finite total curvature, and hence, by Huber's theorem, M is compact with finitely many ends. By Theorem A, each end E of M converges exponentially to a Delaunay surface Σ. Since M is embedded, the Delaunay surface Σ to which an end E converges has to be an onduloid or a right circular cylinder.

We first claim that the Delaunay surface Σ towards which E converges is actually a cylinder.

Suppose it is not. By a rigid motion, we can assume that the axis of Σ is parallel to the y axis and meets the z axis. Then, there is a value z_0 of z such that the line $y \to (0, y, z_0)$ intersects Σ infinitely often. Since E approaches Σ at infinity, the algebraic equation $p(0, y, z_0) = 0$ has infinitely many solutions. This is impossible. So Σ is a cylinder as we claimed.

We claim now that E contains a open set of the cylinder Σ.

To see this, we take a rigid motion so that one of the straight lines of the cylinder Σ agrees with the coordinate y-axis. Thus, one of the intersection curves of E with the plane $x = 0$ is a curve β that converges to the y-axis. If y is large enough, β is given by

$$\beta(y) = (0, y, z(y)),$$

where $z(y)$ is a function that satisfies

$$\lim_{y\to\infty} z(y) = 0$$

Since the curve β belongs to the end E, we have

$$p(0, y, z(y)) = 0 \tag{3}$$

Observe that the polynomial p can be written as

$$p(x, y, z) = a_n z^n + a_{n-1} z^{n-1} + \ldots + a_1 z + a_0$$

where $a_k = a_k(x, y)$ is a polynomial in x and y of degree $\leq n$. By Theorem A, we have that

$$\lim_{y\to\infty} z(y) = \lim_{y\to\infty} Ce^{-\lambda y} = 0$$

By a known result in Calculus, we have, for any integer k,

$$\lim_{y\to\infty} y^k e^{-\lambda y} = 0$$

for any integer k.

Thus, by computing the limit in the equation $p(0, y, z(y)) = 0$ as $y \to \infty$ along the curve β, we obtain that a_0 does not depend on Y, and $a_0 = 0$. This means that, for any y, the equation $p(0, y, z) = 0$ has $z = 0$ as a root, i.e., the straight line $y \to (0, y, 0)$ is contained in E.

The above argument applies to an arbitrary straight line of Σ. It follows that an open set in E is a cylinder. This proves our claim

Thus, there exists an open set U in M with the property that the Gaussian curvature K vanishes in M. Since M is analytic, K vanishes identically in M. It is then well known (see e.g. [8]) that M is a cylinder. Since H is constant, this is a circular cylinder. This proves the Theorem.

Remark: A crucial point in the proof is that the convergence in [7] is exponential. It allows us to prove that not only an arbitrary line in the cylinder Σ converges to E but that actually it is contained in E.

4 Final Remarks

The case $H = 0$. There are many algebraic minimal surfaces in $\mathbb{R}^3$ (see p. 161 of the English translation of Nitsche's book [10]). However, the examples we are most familiar with, namely, the Enneper surface and the Hennenberg surface, are not embedded; thus they are not regular algebraic surfaces.

In fact it is simple to prove the following proposition

Proposition: *There are no regular algebraic minimal surfaces in $\mathbb{R}^3$ except the plane.*

Proof: Let M be an algebraic surface in $\mathbb{R}^3$. As we have seen, such surface it is finitely connected, i.e., it is a compact surface with a finite number of ends. We also know that M is properly embedded.

Let E be one of its ends. Parametrically E can be described by a map $x : D - \{O\} \to \mathbb{R}^3$, where D is an open disk of $\mathbb{R}^2$ centered at the origin and O is the origin.

We may assume, after a rotation if necessary, that the Gauss map, which extends to O (see Osserman [12]), takes on the value $(0, 0, 1)$ at O. The two simplest examples of such ends are the plane and (either end of) the catenoid.

Now we use a result proved by R. Schoen [14]. He showed that such an end is the graph of the function x_3 defined over the (x_1, x_2)-plane and

When $a \neq 0$ the end is of catenoid type. When $a = 0$ the end of the planar type. In fact, if $a \neq 0$ the function x_3 will be asymptotic to the graph of the function $\log \rho$; if $a = 0$ it will be asymptotic to the graph of a constant function (equal to β).

Let's assume that E is of the catenoid type. Consider the curve α intersection of the E with the plane $x_2 = 0$ in the region $x_1 > 0$. Since M is given by the equation $p(x_1, x_2, x_3) = 0$, then the curve α is algebraic, given by $p(x_1, 0, x_3) = 0$. This curve must be asymptotic to the graph of the function $x_3 = a \log x_1$. But this is impossible. Hence, M can not have end of the catenoid type.

Thus, all the ends of M are of the planar type. But they are in finite number. Since M is embedded, the planes asymptotic to M must be parallel. It follows that there are two parallel planes such that M is contained in the region bounded by them. It follows by the halfspace theorem for minimal surfaces [3] that M must be a plane.

Hypersurfaces in $\mathbb{R}^{n+1}$, $n \geq 3$. In this case we consider the zeros of a polynomial function $p(x_0, x_1, \ldots, x_n)$, $n \geq 3$, with $\nabla p \neq 0$ everywhere, and call it *regular algebraic hypersurfaces* M^n of $\mathbb{R}^{n+1}$. Similar to the case $n = 2$, the only compact example of such hypersurfaces are spheres. This follows immediately from Alexandrov theorem. So, we are left to consider the complete non-compact case. A generalized cylinder C^k in $\mathbb{R}^{n+1}$ is a product $B^k \times \mathbb{R}^{n-k}$, where the basis $B^k \subset \mathbb{R}^{k+1} \subset \mathbb{R}^{n+1}$ is a hypersurface of $\mathbb{R}^{k+1}$ and the product is embedded in $\mathbb{R}^{n+1}$ in the canonical way, i.e., $B^k \times \mathbb{R}^{n-k} \subset \mathbb{R}^{k+1} \times \mathbb{R}^{n-k}$. It is easily checked that when B is a k-sphere, C^k has nonzero constant mean curvature. The following lemma is again a consequence of Alexandrov's theorem.

Lemma. *Let C^k be an algebraic regular generalized cylinder in $\mathbb{R}^{n+1}$ whose basis B is a compact hypersurface. If C^k has constant mean curvature then the bases B^k is a k-sphere.*

We do not know any further examples of a regular algebraic hypersurfaces

in $I\!R^{n+1}$, $n > 2$, with nonzero constant mean curvature. We can ask a question similar to the one we answered for $n = 2$. The possible extension of our proof, however, needs new ideas. Although the total Gauss-Kronecker curvature is again finite, there is no Huber theorem for $n > 2$, and the proof of the structure theorem of [7] does not work por hypersurfes in $I\!R^{n+1}$, $n > 2$.

References

[1] Alexandrov, A.D., *A characteristic property of spheres*, Ann. Math. Pura App. **58** (1962) 303-315.

[2] do Carmo, M., *Differential Geometry of Curves and Surfaces*, Prentice Hall, 1976.

[3] Hoffman, D. and Meeks III, W. H., *The strong halfspace theorem for minimal surfaces*. Inventiones Math. **101** (1990)373377.

[4] Hopf, H., *Über Flächen mit einer relation zwischen hauptkrümungen*, Math. Nachr. **4** (1951) 232-249.

[5] Hopf, H., *Differential Geometry in the large*, Springer Lecture Notes in Mathematics **1000**, 2^{nd} edition, 1989.

[6] Huber, A., *On subharmonic functions and differential geometry in the large*, Comment. Math. Helv. **32** (1957) 13-72.

[7] Korevaar, N.J., Kusner, R., and Solomon, B., *The structure of complete embedded surfaces with constant mean curvature*, J. Diff. Geometry **30** (1989) 465-503.

[8] Massey, W.S., *Surfaces of Gaussian curvature zero in Euclidean spaces*, Tohoku Math. J. **14** (1962) 73-79.

[9] Meeks III, W., *The topology and geometry of embedded surfaces of constant mean curvature*, J. Diff. Geometry **27** (1988) 539-552.

[10] Nitsche, J.C.C., Lectures in Minimal surfaces, vol. 1, Cambridge University Press, Cambridge, 1988.

[11] Osserman, R., *The total curvature of algebraic surfaces*, Contributions to Analysis and Geometry, 249-257, Johns Hopkins University Press, Baltimore, 1982.

[12] Osserman, R., *Global properties of minimal surfaces in E^3 and E^n*, Annals of Mathematics **80** (1964) 340-364.

[13] Perdomo, O.M., *Algebraic constant mean curvature surfaces in Euclidean space*, Houston Journal of Mathematics **39** (2013) 127-136.

[14] Schoen, R., *Uniqueness, symmetry, and embeddedness of minimal surfaces*, Journal of Differential Geometry **18** (1983)791809.

[15] Whitney, H., *Geometric Integration Theory*, Princeton University Press, 1957.

Manfredo Perdigão do Carmo
Instituto Nacional de Matemática Pura e Aplicada - IMPA
Estrada Dona Castorina 110
22460-320 Rio de Janeiro - RJ
Brazil
manfredo@impa.br

João Lucas Marques Barbosa
Rua Carolina Sucupira 723 Ap 2002
60140-120 Fortaleza - Ce
Brazil
joaolucasbarbosa@terra.com.br